

ET4350  
Applied Convex Optimization  
Lecture 5

## Unconstrained minimization

$$\text{minimize } f(x)$$

- $f$  convex, twice continuously differentiable (hence  $\text{dom } f$  open)
- we assume optimal value  $p^* = \inf_x f(x)$  is attained (and finite)

### unconstrained minimization methods

- produce sequence of points  $x^{(k)} \in \text{dom } f$ ,  $k = 0, 1, \dots$  with

$$f(x^{(k)}) \rightarrow p^*$$

- can be interpreted as iterative methods for solving optimality condition

$$\nabla f(x^*) = 0$$

## Descent methods

$$x^{(k+1)} = x^{(k)} + t^{(k)} \Delta x^{(k)} \quad \text{with } f(x^{(k+1)}) < f(x^{(k)})$$

- other notations:  $x^+ = x + t\Delta x$ ,  $x := x + t\Delta x$
- $\Delta x$  is the *step*, or *search direction*;  $t$  is the *step size*, or *step length*
- from convexity,  $f(x^+) < f(x)$  implies  $\nabla f(x)^T \Delta x < 0$   
(i.e.,  $\Delta x$  is a *descent direction*)

---

*General descent method.*

**given** a starting point  $x \in \text{dom } f$ .

**repeat**

1. Determine a descent direction  $\Delta x$ .
2. *Line search.* Choose a step size  $t > 0$ .
3. *Update.*  $x := x + t\Delta x$ .

**until** stopping criterion is satisfied.

---

## Line search types


**exact line search:**  $t = \operatorname{argmin}_{t>0} f(x + t\Delta x)$

**backtracking line search** (with parameters  $\alpha \in (0, 1/2)$ ,  $\beta \in (0, 1)$ )

- starting at  $t = 1$ , repeat  $t := \beta t$  until

$$f(x + t\Delta x) < f(x) + \alpha t \nabla f(x)^T \Delta x$$

- graphical interpretation: backtrack until  $t \leq t_0$



# Gradient descent method

general descent method with  $\Delta x = -\nabla f(x)$

---

**given** a starting point  $x \in \text{dom } f$ .

**repeat**

1.  $\Delta x := -\nabla f(x)$ .

2. *Line search.* Choose step size  $t$  via exact or backtracking line search.

3. *Update.*  $x := x + t\Delta x$ .

**until** stopping criterion is satisfied.

---

- stopping criterion usually of the form  $\|\nabla f(x)\|_2 \leq \epsilon$
- convergence result: for strongly convex  $f$ ,

$$f(x^{(k)}) - p^* \leq c^k (f(x^{(0)}) - p^*)$$

$c \in (0, 1)$  depends on  $m$ ,  $x^{(0)}$ , line search type

- very simple, but often very slow; rarely used in practice


## quadratic problem in $\mathbb{R}^2$

$$f(x) = (1/2)(x_1^2 + \gamma x_2^2) \quad (\gamma > 0)$$

with exact line search, starting at  $x^{(0)} = (\gamma, 1)$ :


$$x_1^{(k)} = \gamma \left( \frac{\gamma - 1}{\gamma + 1} \right)^k, \quad x_2^{(k)} = \left( -\frac{\gamma - 1}{\gamma + 1} \right)^k$$

- very slow if  $\gamma \gg 1$  or  $\gamma \ll 1$
- example for  $\gamma = 10$ :




## nonquadratic example

$$f(x_1, x_2) = e^{x_1+3x_2-0.1} + e^{x_1-3x_2-0.1} + e^{-x_1-0.1}$$




backtracking line search



exact line search

a problem in  $\mathbf{R}^{100}$

$$f(x) = c^T x - \sum_{i=1}^{500} \log(b_i - a_i^T x)$$



'linear' convergence, *i.e.*, a straight line on a semilog plot



## Steepest descent method

**normalized steepest descent direction** (at  $x$ , for norm  $\|\cdot\|$ ):

$$\Delta x_{\text{nsd}} = \operatorname{argmin}\{\nabla f(x)^T v \mid \|v\| = 1\}$$

interpretation: for small  $v$ ,  $f(x + v) \approx f(x) + \nabla f(x)^T v$ ;

direction  $\Delta x_{\text{nsd}}$  is unit-norm step with most negative directional derivative

**(unnormalized) steepest descent direction**

$$\Delta x_{\text{sd}} = \|\nabla f(x)\|_* \Delta x_{\text{nsd}}$$

satisfies  $\nabla f(x)^T \Delta x_{\text{sd}} = -\|\nabla f(x)\|_*^2$


**steepest descent method**

- general descent method with  $\Delta x = \Delta x_{\text{sd}}$
- convergence properties similar to gradient descent


## examples

- Euclidean norm:  $\Delta x_{\text{sd}} = -\nabla f(x)$
- quadratic norm  $\|x\|_P = (x^T P x)^{1/2}$  ( $P \in \mathbf{S}_{++}^n$ ):  $\Delta x_{\text{sd}} = -P^{-1} \nabla f(x)$
- $\ell_1$ -norm:  $\Delta x_{\text{sd}} = -(\partial f(x)/\partial x_i) e_i$ , where  $|\partial f(x)/\partial x_i| = \|\nabla f(x)\|_\infty$

unit balls and normalized steepest descent directions for a quadratic norm and the  $\ell_1$ -norm:



## choice of norm for steepest descent



- steepest descent with backtracking line search for two quadratic norms
- ellipses show  $\{x \mid \|x - x^{(k)}\|_P = 1\}$
- equivalent interpretation of steepest descent with quadratic norm  $\|\cdot\|_P$ : gradient descent after change of variables  $\bar{x} = P^{1/2}x$

shows choice of  $P$  has strong effect on speed of convergence

## Newton step

$$\Delta x_{\text{nt}} = -\nabla^2 f(x)^{-1} \nabla f(x)$$


### interpretations

- $x + \Delta x_{\text{nt}}$  minimizes second order approximation

$$\hat{f}(x + v) = f(x) + \nabla f(x)^T v + \frac{1}{2} v^T \nabla^2 f(x) v$$


- $x + \Delta x_{\text{nt}}$  solves linearized optimality condition

$$\nabla f(x + v) \approx \nabla \hat{f}(x + v) = \nabla f(x) + \nabla^2 f(x) v = 0$$



- $\Delta x_{nt}$  is steepest descent direction at  $x$  in local Hessian norm

$$\|u\|_{\nabla^2 f(x)} = (u^T \nabla^2 f(x) u)^{1/2}$$



dashed lines are contour lines of  $f$ ; ellipse is  $\{x + v \mid v^T \nabla^2 f(x) v = 1\}$   
 arrow shows  $-\nabla f(x)$

## Newton's method

---

**given** a starting point  $x \in \text{dom } f$ , tolerance  $\epsilon > 0$ .

**repeat**

1. *Compute the Newton step and decrement.*

$$\Delta x_{\text{nt}} := -\nabla^2 f(x)^{-1} \nabla f(x); \quad \lambda^2 := \nabla f(x)^T \nabla^2 f(x)^{-1} \nabla f(x).$$

2. *Stopping criterion. quit* if  $\lambda^2/2 \leq \epsilon$ .

3. *Line search.* Choose step size  $t$  by backtracking line search.

4. *Update.*  $x := x + t\Delta x_{\text{nt}}$ .

---



affine invariant, *i.e.*, independent of linear changes of coordinates:

Newton iterates for  $\tilde{f}(y) = f(Ty)$  with starting point  $y^{(0)} = T^{-1}x^{(0)}$  are

$$y^{(k)} = T^{-1}x^{(k)}$$

## Examples


example in  $\mathbb{R}^2$  (page 10–9)



- backtracking parameters  $\alpha = 0.1$ ,  $\beta = 0.7$
- converges in only 5 steps
- quadratic local convergence

example in  $\mathbf{R}^{10000}$  (with sparse  $a_i$ )

$$f(x) = - \sum_{i=1}^{10000} \log(1 - x_i^2) - \sum_{i=1}^{100000} \log(b_i - a_i^T x)$$



- backtracking parameters  $\alpha = 0.01$ ,  $\beta = 0.5$ .
- performance similar as for small examples